


## Intellectual Property Network for Business

IPN Home | Search | Order | Shopping Cart | Login | Help



## WO9709049A2: METHOD FOR EFFECTING VASODILATION WITH (1,5-INTER)ARYL PROSTAGLANDIN DERIVATIVES

[View Images \(31 pages\)](#) | [View Cart](#)
[Premium Data 1: PDF \(~2950 KB\)](#) | [TIFF \(~2330 KB\)](#) | [Fax](#) | [More choices...](#)

**Inventor(s):** CHEN, June , 28771 Via Los Arboles, San Juan Capistrano, CA 92675, United States of America  
 BURK, Robert, M. , 1337 Cerritos Drive, Laguna Beach, CA 92651, United States of America  
 WOODWARD, David, F. , 23152 Tulip, Lake Forest, CA 92630, United States of America

**Applicant(s):** ALLERGAN, 8301 Mars Drive, Waco, TX 76712, United States of America

**Issued/Filed Dates:** March 13, 1997 / Aug. 28, 1996

**Application Number:** WO1996US0013889

**IPC Class:** A61K 031/557; C07C 405/00;

**Priority Number(s):** Sept. 1, 1995 US1995008522775

**Designated Countries:** AU, CA, JP, European patent: AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE





**Abstract:** The present invention provides a method of effecting vasodilation, comprising: administering to a warm blooded animal in need of such treatment, an effective amount of (1,5-inter)aryl prostaglandin derivative represented by formula (I), wherein n is 0 or an integer of from 1 to 6, R is selected from the group of radicals represented by the formulae CO<sub>2</sub>R', CONR'<sub>2</sub>, CH<sub>2</sub>OR' and SO<sub>2</sub>NR'<sub>2</sub>', wherein R' is hydrogen or a lower alkyl radical having from one to six carbon atoms; R'' is hydrogen or an acyl radical having the formula: (CO)R''' wherein R''' is a saturated or unsaturated acyclic hydrocarbon radical having from 1 to about 10 carbon atoms, or -(CH<sub>2</sub>)<sub>m</sub>R'''' wherein m is 0 or an integer of from 1 to 6 and R'''' is an aliphatic ring having from 3 to 7 carbon atoms or an aryl group, e.g. phenyl, or a heteroaryl group, e.g. thienyl, furanyl or pyridyl, and preferably R''' is a lower alkyl group having from 1 to 6 carbon atoms; the hatched triangular segments represent alpha oriented bonds, the solid triangular segments represent beta oriented bonds and the wavy segments represent bonds that may be in either the cis or trans orientation. More preferably said (1,5 inter)aryl prostaglandin derivative is a compound represented by formula (II). Most preferably, said (1,5-inter)aryl prostaglandin derivative is a compound of formula (II) wherein R is CO<sub>2</sub>R', R' is hydrogen and n is 0.

[Show "fr" Abstract]

Best Available Copy

# Best Available Copy

Representative  
Image:



[Show "fr" image]

Attorney, Agent, or  
Firm  
Foreign References:

BARAN, Robert, J.;

none

(No patents reference this one)